

Ovarian Cancer Symptoms

Traditionally Ovarian Cancer was known as ‘the Silent Killer’, however recent research* shows the following symptoms are found more frequently in women with ovarian cancer:

Key Diagnostic Symptoms

- Bloating
- Pelvic or abdominal pain
- Difficulty eating or feeling full quickly
- Urinary symptoms (urgency or frequency)

When these symptoms persist for more than a few weeks or represent a change from *your* normal, see your doctor – preferably a gynecologist.

The frequency and/or number of symptoms are keys in the diagnosis of ovarian cancer. Even early stage ovarian cancer can produce

Other Common Symptoms

- Fatigue
- Indigestion
- Back pain
- Pain with intercourse
- Constipation

While common, these other symptoms are less diagnostically helpful since they are also found in many women *without* ovarian cancer.

Provided by www.ovca.net

* Goff BA, Mandel L, Drescher CW, Urban N, Gough S, Schurman K, Patras JJ, Mahony BS, Anderson M *DeveLopment of an ovarian cancer symptom index*. Cancer 2007; 109:221-7. Level II-2

Ovarian Cancer Symptoms

Traditionally Ovarian Cancer was known as ‘the Silent Killer’, however recent research* shows the following symptoms are found more frequently in women with ovarian cancer:

Key Diagnostic Symptoms

- Bloating
- Pelvic or abdominal pain
- Difficulty eating or feeling full quickly
- Urinary symptoms (urgency or frequency)

When these symptoms persist for more than a few weeks or represent a change from *your* normal, see your doctor – preferably a gynecologist.

The frequency and/or number of symptoms are keys in the diagnosis of ovarian cancer. Even early stage ovarian cancer can produce

Other Common Symptoms

- Fatigue
- Indigestion
- Back pain
- Pain with intercourse
- Constipation

While common, these other symptoms are less diagnostically helpful since they are also found in many women *without* ovarian cancer.

Provided by www.ovca.net

* Goff BA, Mandel L, Drescher CW, Urban N, Gough S, Schurman K, Patras JJ, Mahony BS, Anderson M *DeveLopment of an ovarian cancer symptom index*. Cancer 2007; 109:221-7. Level II-2

Ovarian Cancer Symptoms

Traditionally Ovarian Cancer was known as ‘the Silent Killer’, however recent research* shows the following symptoms are found more frequently in women with ovarian cancer:

Key Diagnostic Symptoms

- Bloating
- Pelvic or abdominal pain
- Difficulty eating or feeling full quickly
- Urinary symptoms (urgency or frequency)

When these symptoms persist for more than a few weeks or represent a change from *your* normal, see your doctor – preferably a gynecologist.

The frequency and/or number of symptoms are keys in the diagnosis of ovarian cancer. Even early stage ovarian cancer can produce

Other Common Symptoms

- Fatigue
- Indigestion
- Back pain
- Pain with intercourse
- Constipation

While common, these other symptoms are less diagnostically helpful since they are also found in many women *without* ovarian cancer.

Provided by www.ovca.net

* Goff BA, Mandel L, Drescher CW, Urban N, Gough S, Schurman K, Patras JJ, Mahony BS, Anderson M *DeveLopment of an ovarian cancer symptom index*. Cancer 2007; 109:221-7. Level II-2

Ovarian Cancer Symptoms

Traditionally Ovarian Cancer was known as ‘the Silent Killer’, however recent research* shows the following symptoms are found more frequently in women with ovarian cancer:

Key Diagnostic Symptoms

- Bloating
- Pelvic or abdominal pain
- Difficulty eating or feeling full quickly
- Urinary symptoms (urgency or frequency)

When these symptoms persist for more than a few weeks or represent a change from *your* normal, see your doctor – preferably a gynecologist.

The frequency and/or number of symptoms are keys in the diagnosis of ovarian cancer. Even early stage ovarian cancer can produce

Other Common Symptoms

- Fatigue
- Indigestion
- Back pain
- Pain with intercourse
- Constipation

While common, these other symptoms are less diagnostically helpful since they are also found in many women *without* ovarian cancer.

Provided by www.ovca.net

* Goff BA, Mandel L, Drescher CW, Urban N, Gough S, Schurman K, Patras JJ, Mahony BS, Anderson M *DeveLopment of an ovarian cancer symptom index*. Cancer 2007; 109:221-7. Level II-2